

Primordial Non-Gaussianity from LAMOST Surveys

Yan Gong^{1,2}, Xin Wang^{1,2}, Zheng Zheng³ and Xuelei Chen^{1,4,5} *

¹National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, China

²Graduate School of Chinese Academy of Sciences, Beijing 100049, China

³Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA

⁴Center of High Energy Physics, Peking University, Beijing 100871, China

⁵Kavli Institute for Theoretical Physics China, CAS, Beijing 100190, China

Abstract The primordial non-Gaussianity (PNG) in matter density perturbation is a very powerful probe of the physics of the very early Universe. The local PNG can induce a distinct scale-dependent bias on the large scale structure distribution of galaxies and quasars, which could be used for constraining it. We study the detection limits on PNG from the surveys of the LAMOST telescope. The cases of the main galaxy survey, the luminous red galaxy (LRG) survey, and the quasar survey of different magnitude limits are considered. We find that the MAIN1 sample (i.e. the main galaxy survey with one magnitude deeper than the SDSS main galaxy survey, or $r < 18.8$) could only provide very weak constraint on PNG. For the MAIN2 sample ($r < 19.8$) and the LRG survey, the $2\sigma(95.5\%)$ limit on the PNG parameter f_{NL} are $|f_{\text{NL}}| < 145$ and $|f_{\text{NL}}| < 114$ respectively, comparable to the current limit from cosmic microwave background (CMB) data. The quasar survey could provide much more stringent constraint, and we find that the 2σ limit for $|f_{\text{NL}}|$ is between 50 and 103, depending on the magnitude limit of the survey. With Planck-like priors on cosmological parameters, the quasar survey with $g < 21.65$ would improve the constraints to $|f_{\text{NL}}| < 43$ (2σ). We also discuss the possibility of further tightening the constraint by using the relative bias method proposed by Seljak (2008).

Key words: cosmology:large scale structure

preprint number: CAS-KITPC/ITP-161, arXiv:0904.4257

1 INTRODUCTION

The seed of large scale structure (LSS) is usually thought to be produced during a period of inflation in the early universe. Models of the standard scenario of single field slow roll inflation generally predict that these primordial perturbations are Gaussian distributed (Maldacena, 2003; Acquaviva et al., 2003; Creminelli, 2003; Bartolo, Komatsu, & Matarrese, 2004). However, significant primordial non-Gaussianity (PNG) may be produced by multi-field inflation (Linde & Mukhanov, 1997) and reheating (Dvali, Gruzinov, & Zaldarriaga, 2004), if the inflaton potential has some sharp features (Chen, Easther, & Lim, 2007), or if the inflation is driven by field with non-canonical kinetic terms (Chen, Huang, Kachru, Shiu, 2007). PNG is also predicted by alternative models to inflation, such as the ekpyrotic scenario (Lehners & Steinhardt, 2008). Thus, non-Gaussianity in primordial density perturbations could be a very powerful discriminator for the physical models of the very early Universe (Bartolo, Komatsu, & Matarrese, 2004; Komatsu et al., 2009).

* E-mail: xuelei@cosmology.bao.ac.cn

The PNG in density perturbation is usually parametrized as

$$\Phi(\mathbf{x}) = \phi(\mathbf{x}) + f_{\text{NL}}(\phi^2(\mathbf{x}) - \langle \phi^2 \rangle), \quad (1)$$

where Φ is Bardeen's gauge-invariant potential and ϕ is a Gaussian random field. The parameter f_{NL} , which denotes the effect of the PNG, is typically of order 10^{-2} in standard inflation models (e.g., Gangui et al. 1994). Non-linear transformation from primordial field fluctuation to observables generically gives rise to $f_{\text{NL}} \sim O(1)$. With $\phi \sim 10^{-5}$, this produces an extremely small non-Gaussianity in observables, such as the cosmic microwave background (CMB) temperature anisotropy or LSS bispectrum.

The CMB anisotropy so far offers the cleanest observational probe of PNG (Wang & Kamionkowski, 2000; Verde et al., 2001). Yadav & Wandelt (2008) recently claimed detection of a positive f_{NL} at 99.5% significance ($27 < f_{\text{NL}} < 147$ at 2σ) from the Wilkinson Microwave Anisotropy Probe (WMAP) 3-year data. Analysis of the WMAP 5-year data using the bispectrum also favors a slightly positive f_{NL} , $-9 < f_{\text{NL}} < 111$ at 95.5% confidence level (C.L.) (Komatsu et al., 2008), while the Minkowski functional method yields $-178 < f_{\text{NL}} < 64$ (Komatsu et al., 2008). In the future, the Planck satellite should significantly improve the constraint on f_{NL} , which may eventually push the uncertainty in f_{NL} to be around 5 (Komatsu & Spergel, 2001). Alternatively, one may use the LSS observations to search for the PNG. However, even if the initial fluctuations are Gaussian, during the gravitational growth of the LSS, non-linear effect also generates non-Gaussianity, which may swamp the primordial one. Typically, the LSS constraints on PNG using the bispectrum technique are weak (Verde et al., 2001; Scoccimarro, Sefusatti & Zaldarriaga, 2004; Sefusatti & Komatsu, 2007). One consequence of PNG in structure formation is to significantly enhance the number of the rare peaks of fluctuation, boosting the formation of massive objects (Lucchin, Matarrese & Vittorio, 1986; Matarrese, Verde & Jimenez, 2000). One could observe this effect from the abundance of galaxy clusters (Benson, Reichardt, & Kamionkowski, 2002) or dark matter halos harboring the first generation of stars (Chen et al., 2003). However, the required observational task is difficult due to low-number statistics.

Recently, Dalal et al. (2008) proposed a powerful and practical method to probe the local PNG with LSS observations. The PNG-caused enhancement on the formation of massive dark matter halos induces a distinctive scale-dependent bias on the largest scales. Subsequent works confirmed and extended this result (Matarrese & Verde, 2008; Slosar et al., 2008; Afshordi & Tolley, 2008; McDonald, 2008; Carbone, Verde, & Matarrese, 2008; Kamionkowski, Verde, & Jimenez, 2008), showing that useful limits on PNG could be obtained from LSS surveys.

In the present work, we investigate the sensitivity to PNG from the extragalactic surveys of the Large sky Area Multi-Object Spectroscopic Telescope (LAMOST)¹. The LAMOST is a recently built Chinese 4-meter Schmidt telescope with a field of view of 20 deg^2 . Equipped with 4000 optical fibers that are individually positioned by computer during observation, it can take spectra of 4000 targets simultaneously (Chu, 1998; Stone, 2008). The LAMOST is currently undergoing engineering tests and calibrations, and it is expected to begin conducting surveys by 2010/2011. In a recent paper, we have forecasted the constraining power of the LAMOST extragalactic surveys on dark energy measurement (Wang et al., 2008) by considering different samples of galaxies and quasars. Here, we make similar forecast on the constraining power on PNG.

2 METHOD

On large scales, the PNG induces a distinct scale-dependent bias in the distribution of tracer objects, which can be measured from the correlation function or power spectrum of galaxies and quasars. Extending the work of Dalal et al. (2008), Slosar et al. (2008) found that this bias departs from the Gaussian case as

$$\Delta b(z, M, k) = 2f_{\text{NL}}(b - p)\delta_c \frac{3\Omega_{m0}}{2k^2 T(k)D(z)} \left(\frac{H_0}{c}\right)^2, \quad (2)$$

¹ <http://www.lamost.org/>

where b is the Eulerian bias and can be calculated as usual with the halo model, $\delta_c = 1.686$, $D(z)$ is the linear growth factor normalized to be the scale factor a in the matter domination epoch, Ω_{m0} is the present matter energy density parameter, $T(k)$ is the transfer function, H_0 is the current Hubble parameter and c is the speed of light.

The parameter p is introduced to account for the effect of merger bias. If the effect of merger on the formation of tracer objects is negligible, its value should be unity. However, the formation of some objects may be triggered by mergers. For example, quasars are believed to be accreting massive black holes in the center of galaxies, and the activity of quasar may be triggered by galaxy mergers, which send gas to the center of the galaxy, providing the materials needed for accretion. If the merger effect is dominant, $p = 1.6$. In general, the value of p is between 1 and 1.6.

We consider three types of extragalactic surveys for LAMOST. (1) The main galaxy survey, which is a magnitude-limited general survey of all types of galaxies. By MAIN1 and MAIN2, we denote respectively the samples which are 1 and 2 magnitude deeper than the SDSS surveys, i.e. MAIN1 for $r < 18.8$, and MAIN2 for $r < 19.8$. (2) The LRG survey, a survey of color-selected luminous red galaxies (LRGs). Here for illustration we have used the MegaZ LRG sample (Collister et al., 2006) for calculation, though in reality the LAMOST LRG will be selected independently. (3) The quasar survey. Depending on the magnitude limit, we define three samples, QSO1 ($g < 20.5$), QSO2 ($g < 21$), and QSO3 ($g < 21.65$). As in Wang et al. (2008), we assume that the surveyed area is 8000 deg^2 , for which we have the SDSS imaging data, even though the actual survey may cover a larger area if input catalogue for the south Galactic cap is available. For more details of these samples, including their number densities, redshift distributions, and the values of bias (estimated in the Gaussian models), see Wang et al. (2008).

We model the galaxy power spectrum as

$$P_g(k) = (b_g + \Delta b)^2 P_{\text{lin}}(k) \frac{1 + Qk^2}{1 + Ak^2}, \quad (3)$$

where b_g is the bias of the galaxies calculated for the Gaussian case, P_{lin} is the linear matter power spectrum (Eisenstein & Hu, 1998), k is the wavenumber ($10^{-2} \text{ hMpc}^{-1} < k < 0.2 \text{ hMpc}^{-1}$), Q and A are free parameters which describe the non-linear effect. For fiducial models, we fix $Q = 4.6$ for the main samples, $Q = 30$ for the LRG sample, and adopt $A = 1.4$ (Cole et al., 2005; Tegmark et al., 2006). For the quasar power spectrum, we adopt the simpler form

$$P_q(k) = (b_q + \Delta b)^2 P_{\text{lin}}(k), \quad (4)$$

where b_q is the bias of quasars. The value of bias for each of the sample has been given in Wang et al. (2008). We assume that the PNG is small, so that the value of bias does not change much.

The deviation of bias could be discovered with precision measurement of the power spectrum of the galaxy or quasar distribution. The measurement error of the power spectrum is given by (Feldman, Kaiser, Peacock, 1994; Tegmark, 1997)

$$\frac{\sigma_P}{P} = 2\pi \sqrt{\frac{1}{V_s k^2 \Delta k} \frac{1 + nP}{nP}} = 2\pi \sqrt{\frac{1}{V_{\text{eff}} k^2 \Delta k}}, \quad (5)$$

where V_s is the survey volume and V_{eff} is the effective volume, defined by

$$V_{\text{eff}} \equiv \int \left[\frac{n(\mathbf{r})P}{n(\mathbf{r})P + 1} \right]^2 d\mathbf{r} = \left[\frac{nP}{nP + 1} \right]^2 V_s. \quad (6)$$

The second equality holds if the comoving number density $n(\mathbf{r})$ is a constant. We divide the samples into different redshift bins as in Wang et al. (2008), then the total likelihood function is the product of the likelihood function for all the different bins.

In order to estimate the error on the measurement of f_{NL} , we use a Markov Chain Monte Carlo (MCMC) technique to explore the parameters space, then marginalize over (i.e., integrate out) the other




Fig. 1 Fiducial values of bias and the fits with the functional form $b(z) = b_0(1+z)^\gamma$ for the Main1, LRG, and QSO3 samples. For each sample, the fiducial bias values (symbols) are from Wang et al. (2008), the thick curve are the best fit with the thin curves denotes the central 68% distribution marginalized over the MCMC results.

parameters to obtain a constraint on the PNG. For details of the MCMC technique, see, e.g., our earlier paper (Gong & Chen, 2007). The fiducial cosmology we adopt is the WMAP 5-year best fit Λ CDM model without PNG, with $\Omega_{b0} = 0.05$, $\Omega_{m0} = 0.24$, $\Omega_{\Lambda0} = 0.76$, $\sigma_8 = 0.82$, $n_s = 0.96$, $f_{NL} = 0.0$ and $h_0 = 0.7$. The allowed ranges of parameters in the Monte Carlo are $\Omega_{b0} \in (0, 0.1)$, $\Omega_{m0} \in (0, 1)$, $\sigma_8 \in (0.6, 0.9)$, $n_s \in (0.9, 1.1)$, $f_{NL} \in (-400, 400)$, $Q \in (0, 40)$ and $h_0 \in (0.5, 0.9)$.

The value of bias for the sample is an important parameter which is partially degenerate with f_{NL} . We model the value of bias at different redshifts bins in two possible forms:

$$b(z) = b_0(1+z)^\gamma, \quad (7)$$

or

$$b(z) = c_1 + c_2/D_0(z) \quad (8)$$

where $D_0(z)$ is the growth factor normalized such that $D_0(z=0) = 1$. This latter form is inspired by $b(z) = 1 + (b_p - 1)/D_0(z)$, which can be derived in the linear regime for a passively evolved tracer population without any mergers by using $\dot{\delta}_g = \dot{\delta}_m$ and $\delta_g(z) = b(z)\delta_m(z)$ (Fry, 1996). Here b_p is the present day bias factor of the tracer, δ_g and δ_m are the tracer and matter density contrast, respectively. We then try to use these two parametrization forms to fit the bias for our samples which were given in Wang et al. (2008). Our test show that for the second form (Eq.8) to yield good fit, we must have $c_1 < 0$, which is physically unphysical given the original motivation for this form of bias evolution. Thus, we choose to use the first form (Eq. 7) to fit for biases at different redshifts. The parameter b_0 and γ are then free parameters in the MCMC with the priors given by $b_0 \in (0, 10)$ and $\gamma \in (-5, 5)$. The predicted values of bias given by the best fit parameter for several samples (Main1, LRG, QSO3) and those given by Wang et al. (2008) are shown in Fig. 1. We see that the functional form we adopt (Eq.7) is adequate to describe the value of bias as a function of redshift for each sample.

Note that to see how the LSS probes the PNG, here we do not consider joint constraint on f_{NL} from CMB observations (e.g., bispectrum). We also do not use the currently available CMB angular power spectrum data in our MCMC unless otherwise stated (see next section about Planck prior).

3 RESULTS AND CONCLUSIONS

The result of our analysis, i.e., the 1σ (68.3%), 2σ (95.5%), 3σ (99.7%) limits of f_{NL} inferred from the power spectrum of the galaxy and quasar samples are given in Table 1 and Table 2, and also shown in Fig. 2. We see that for the MAIN1 sample, the 3σ (99.7%) limit on $|f_{\text{NL}}|$ reaches 379, so it provides only very weak constraint on PNG. The MAIN2 and LRGs surveys have almost the same effective volume on large scales (Wang et al., 2008), so they provide comparable constraint on f_{NL} . Comparing with the current SDSS spectroscopic LRG survey (Slosar et al., 2008), the constraint on $|f_{\text{NL}}|$ could be improved by a factor of about 1.5, to reach a 2σ limit of about 114, which is comparable to current limits from WMAP data (Komatsu et al., 2008).

Quasars are luminous and can be observed at high redshifts, so one could obtain large survey volume with quasars. For the large scales considered here, the quasars provide the largest effective volume (c.f. Fig.3 of Wang et al. 2008), so we expect the LAMOST quasar surveys to provide a better constraint on f_{NL} . To account for the possible effect of mergers, we calculated for the case of $p = 1$ (non-merger) and $p = 1.6$ (merger). We see that even for the QSO1 sample ($g < 20.5$), the 95.5%(2σ) limit on $|f_{\text{NL}}|$ reaches 103 for the merger case and 82 for the non-merger case. For the QSO2 sample ($g < 21$), the corresponding limits are 82(merger) and 63 (non-merger). For the QSO3 sample ($g < 21.65$), the limits shrink to 69 (merger) and 50 (non-merger). Comparing with the SDSS quasar surveys (Slosar et al., 2008), the LAMOST could improve the constraint of f_{NL} by a factor of 1.3. We note that we assumed a lower QSO bias than Slosar et al. (2008), so the limits obtained here are conservative.

Table 1 The 1σ , 2σ , and 3σ range of f_{NL} constrained with the LAMOST Galaxy Surveys. The last row (LRG- α) is from the relative bias method proposed by Seljak (2008).

Survey	1σ	2σ	3σ
Main1	± 127	± 252	± 379
Main2	± 71	± 145	± 206
LRG	± 53	± 114	± 153
LRG- α	± 13	± 28	± 37

Table 2 The 1σ , 2σ , and 3σ range of f_{NL} constrained with the LAMOST Quasar Surveys.

Survey	Merger ($p = 1.6$)			Non-merger ($p = 1$)		
	1σ	2σ	3σ	1σ	2σ	3σ
QSO1	± 52	± 103	± 155	± 40	± 82	± 122
QSO2	± 39	± 82	± 118	± 31	± 63	± 95
QSO3	± 33	± 69	± 101	± 25	± 50	± 75

The posterior probability distribution function of f_{NL} appears to be nearly Gaussian here, so the 1σ , 2σ , and 3σ errors are almost uniformly spaced. We have also investigated the degeneracy between f_{NL} and Ω_{m0} and n_s . We find that f_{NL} is positively correlated with Ω_{m0} , but it is almost independent with n_s , confirming the results of Slosar et al. (2008).

By the time that the LAMOST surveys are completed, the Planck satellite should have been taken CMB for a few years. With these CMB data, the uncertainties on cosmological parameters should shrink further. Here we simulate this effect by adding priors on the other cosmological parameters based on the forecast of the Planck surveys. We set the standard deviation as $\Omega_{b0} = 0.05 \pm 4.0 \times 10^{-4}$, $\Omega_{m0} = 0.24 \pm 5.0 \times 10^{-3}$, $\sigma_8 = 0.82 \pm 6.7 \times 10^{-3}$, $n_s = 0.96 \pm 4.0 \times 10^{-3}$ and $h_0 = 0.7 \pm 6.9 \times 10^{-3}$ (Colombo et al., 2008). We present the results for the LRG survey and QSO3 survey, and the impacts on other cases are similar. With the stronger priors, the constraints are improved slightly. With the QSO3 sample, we obtain $|f_{\text{NL}}| < 65$ (merger case) and $|f_{\text{NL}}| < 43$ (non-merger case) at 95.5% C.L., in

Table 3 The 1σ , 2σ , and 3σ constraints on f_{NL} from the LAMOST surveys with Planck priors.

Survey	1σ	2σ	3σ
LRG	± 34	± 69	± 102
QSO3 merger	± 32	± 65	± 93
QSO3 non-merger	± 21	± 43	± 60




Fig. 2 The 1σ , 2σ , and 3σ range of f_{NL} from LAMOST surveys. The samples are MAIN1 ($r < 18.8$), MAIN2 ($r < 19.8$), LRG (the MegaZ sample), LRG- α (the constraint with the relative bias method using the LRG and the photometric samples), LRG α -r (the degraded relative bias method considering bias evolution), LRG+P (LRG with Planck priors), QSO1 (QSO with $g < 20.5$, non-merger case), QSO2 (QSO with $g < 21$, non-merger case), QSO3 (QSO with $g < 21.65$, non-merger case), QSO3+P (QSO3 sample, non-merger case, with Planck priors), QSO1-m (QSO1 sample, merger case), QSO2-m (QSO2 sample, merger case), QSO3-m (QSO3 sample, merger case), QSO3-m+P (QSO3 sample, merger case, with Planck priors). The assumed survey area is 8000 deg^2 .

comparison with $|f_{\text{NL}}| < 69$ and $|f_{\text{NL}}| < 50$ without the Planck priors. These results are given in Table 3 and also shown in Fig.2.

Comparing directly the density fields of tracers with different biases in the same survey volume helps to eliminate the measurement error due to cosmic variance and break the degeneracy with other cosmological parameters (Seljak, 2008; McDonald & Seljak, 2008). The relative bias $\alpha = b_1/b_2$ is proposed to be used as the estimator of PNG (Seljak, 2008; McDonald & Seljak, 2008). Under the assumption that the two tracers have no stochasticity in them, and one tracer is unbiased ($b=1$), the constraint of f_{NL} would be improved by a factor of $(2[(\bar{n}_1 P_1)^{-1} + (\bar{n}_2 P_2)^{-1}])^{-1/2}$. The feasibility of this method depends on the number density of the tracers. Here we consider the LRG as the high bias

tracer, and a sample of galaxies with photometric redshift as the low bias tracer. The average redshift of the LRGs is $z \approx 0.4$. Using the halo model method described in Wang et al. (2008), we find that for a sample of galaxies with good photometry up to $r < 21.5$, the average comoving density is $n = 8.11 \times 10^{-3} h^3 \text{Mpc}^{-3}$, and the average bias is $b = 1.08$. This sample of galaxies satisfies the requirement for the low bias tracer and has available SDSS photometry. With the relative bias method, we find that the limit on f_{NL} obtainable with LRGs could be improved by a factor of about 4.1, with the 1σ , 2σ , and 3σ limits on $|f_{\text{NL}}|$ being 13, 28, and 37, respectively.

In fact, since the evolution of the number density and clustering of the two galaxy samples could be different, the assumption of perfect correlation between the two tracer populations may not be valid, especially for the large volumes used for the non-Gaussianity test discussed here. This could introduce an additional error. The detailed study of this effect is beyond the scope of the present paper, we plan to investigate it in subsequent studies. For the problem discussed here, crude estimates show that in the worst case, it may cause a degradation of the statistical power by as large as 50%. The LRG α -r in Fig. 2 shows the degraded precision.

In summary, we forecast the constraints on the primordial non-Gaussianity from the LAMOST surveys. Using the scale-dependent bias effect proposed by Dalal et al. (2008), and based on the estimations of the galaxy and quasar surveys for the LAMOST (Wang et al., 2008), we find that the galaxy surveys provide relatively weak constraint on the f_{NL} parameter, but the QSO surveys could provide strong constraints, with a 2σ limit on $|f_{\text{NL}}|$ somewhere between 50 to 103, depending on the sample size of the LAMOST QSO survey. After adding the Planck priors for other cosmological parameters, the QSO3 survey could give a 2σ limit of 43 for $|f_{\text{NL}}|$. We also considered using the relative bias method (Seljak, 2008). Correlating the LAMOST LRG sample with a photometric sample of galaxies with $r < 21.5$, for which $b \approx 1$ at $z = 0.4$, we obtain a 2σ limit of $|f_{\text{NL}}| < 28$. These results show that the LAMOST could make useful constraints on the primordial non-Gaussianity.

Acknowledgements We thank Anže Slosar for discussions on details of their calculation, and Pengjie Zhang for helpful suggestion. Our MCMC chain computation was performed on the Supercomputing Center of the Chinese Academy of Sciences and the Shanghai Supercomputing Center. This work is supported by the National Science Foundation of China under the Distinguished Young Scholar Grant 10525314, the Key Project Grant 10533010, by the Chinese Academy of Sciences under the grant KJCX3-SYW-N2, and by the Ministry of Science and Technology national basic science Program (Project 973) under grant No. 2007CB815401. Z. Z. gratefully acknowledges support from the Institute for Advanced Study through a John Bahcall Fellowship.

References

- Acquaviva V., Bartolo N., Matarrese S., Riotto, 2003, Nucl. Phys. B667, 119
- Afshordi N., Tolley A. J., 2008, arXiv:0806.1046
- Bartolo, N., Komatsu, E., Matarrese, S., 2004, Phys.Rept., 402, 103.
- Benson, A., Reichardt, C., Kamionkowski, M., 2002, MNRAS, 331, 71.
- Carbone, C., Verde, L., Matarrese, S., 2008, ApJ, 684, L1.
- Chen, X., Cooray, A., Yoshida, N., Sugiyama, N., 2003, MNRAS, 346, L31.
- Chen, X., Easter, R., & Lim, E. A., 2007, JCAP, 0706:023.
- Chen, X., Huang, M., Kachru, S., Shui, G., 2007, JCAP0701:002.
- Chu, Y., 1998, Highlights of Astronomy, 11A., 493.
- Cole S. et al., 2005, MNRAS, 362, 505
- Collister A. et al., 2007, MNRAS 375, 68.
- Colombo L.P.L., Pierpaoli E., Pritchard J.R., 2008, arXiv:0811.2622
- Creminelli P., 2003, JCAP, 0310, 003
- Dalal N., Doré O., Huterer D., Shirokov A., 2008, Phys. Rev. D, 77, 123514
- Dvali, G., Gruzinov, A., Zaldarriaga, M., 2004, Phys. Rev. D, 69, 023505
- Eisenstein D. J., Hu W., 1998, ApJ, 496, 605

- Feldman H. A., Kaiser N., Peacock J. A., 1994 ApJ, 426, 23
Fry, J. N. 1996, ApJ, 461, L65
Gangui, A., Lucchin, F., Matarrese, S., & Mollerach, S. 1994, ApJ, 430, 447
Gong Y., Chen X., 2007, Phys. Rev. D, 76, 123007
Hikage C. et al., 2008, MNRAS, 389, 1439
Kamionkowski, M., Verde, L., & Jimenez, R., 2008, arXiv:0809.0506.
Komatsu E., Spergel D. N., 2001, Phys. Rev. D, 63, 063002
Komatsu, E. et al., 2003, ApJS, 148, 119
Komatsu, E. et al., 2008, arXiv:0803.0547
Komatsu, E. et al., 2009, arXiv:0902.4759, science white paper submitted to the cosmology and fundamental physics science frontier panel of Astro 2010.
Lehners, J.-L., Steinhardt, P.J., 2008, Phys. Rev. D, 78, 023506
Linde A., Mukhanov V., 1997, Phys. Rev. D, 56, 535
Lucchin, F., Matarrese, S., Vittorio, 1986, A&A, 162,13.
Maldacena J., 2003, Journal of High Energy, 5, 13
Matarrese S., Verde L., Jimenez, R., 2000, ApJ, 541, 10.
Matarrese S., Verde L., 2008, ApJ, 667, 77
McDonald P., 2008, arXiv:0806.1061
McDonald P., Seljak U., 2008, arXiv:0810.0323
Scoccimarro, R., Sefusatti, E., Zaldarriaga, M., 2004, Phys. Rev. D, 69, 103513
Sefusatti, E. & Komatsu, E., 2007, Phys. Rev. D, 76, 083004.
Seljak U., 2008, arXiv:0807.1770
Slosar A., Hirata C., Seljak U., Ho S., Padmanabhan N., 2008, JCAP, 08, 031
Stone, R., 2008, Science, 320, 34.
Tegmark M., 1997, Phys. Rev. Lett., 79, 3806
Tegmark M. et al., 2006, Phys. Rev. D, 74, 123507
Verde, L., Jimenez, R., Kamionkowski, M., Matarrese, S., 2001, MNRAS, 325, 412.
Wang, L., Kamionkowski, M., 2000, Phys. Rev. D, 61, 063504.
Wang X., Chen X., Zheng Z., Wu F., Zhang P., Zhao Y., 2009, MNRAS, 394, 1775.
Yadav, A.P.S., Wandelt, B.D., 2008, Phys. Rev. Lett., 100, 181301.